

On factorisable, not simple groups.

— To Professor L. RÉDEI on his 50th birthday.

By J. SZÉP in Szeged.

A group \mathcal{G} is called factorisable by its subgroups \mathcal{H} and \mathcal{K} if each element of \mathcal{G} may be written in the form $G = HK$ with H in \mathcal{H} , K in \mathcal{K} (written $\mathcal{G} = \mathcal{H}\mathcal{K}$).

It was G. ZAPPA [9] who has considered first factorisable groups; he has supposed that the two subgroups have no element in common except the unit element. Since that several papers have been published dealing with factorisable groups [1]—[8]. The general case where the factors may have a subgroup $\mathcal{D} \neq E$ of \mathcal{G} in common was treated by the author and L. RÉDEI [8]. For the case that \mathcal{D} is a normal subgroup G. CASADIO [1] has obtained a similar result as G. ZAPPA.

The interest of factorisable groups may be judged from the fact that an infinite number of simple groups are factorisable [6]. Moreover, it is possible to obtain some criteria for the simplicity of factorisable groups [7].

The present paper contains two principal results:

Theorem 1. *If a group \mathcal{G} of finite order is factorisable by two abelian groups, then \mathcal{G} is not simple.*

Theorem 2. *If a group \mathcal{G} of finite order is factorisable by two abelian groups \mathcal{H} and \mathcal{K} and the orders of \mathcal{H} and \mathcal{K} are relatively prime, then \mathcal{G} is solvable.*

Theorem 2 is connected with the following theorem of O. HÖLDER:

If \mathcal{G} is a group of finite order and if all Sylow-groups of \mathcal{G} are cyclic, then \mathcal{G} is solvable and factorisable by two cyclic groups with relatively prime orders.

The converse is also true, it follows from the original theorem:

If a group \mathcal{G} of finite order is factorisable by two cyclic groups whose orders are relatively prime, then \mathcal{G} is solvable.

Theorem 2 is a generalization of the converse of HÖLDER's theorem.

For the proof of Theorems 1 and 2 it is necessary to recall some facts contained in earlier papers:-

Theorem A. [8] *Let the group \mathfrak{G} be factorisable by the proper subgroups \mathfrak{H} and \mathfrak{K} , and let $\mathfrak{H} \cap \mathfrak{K} = \mathfrak{D} \neq E$. If \mathfrak{D} has a proper subgroup \mathfrak{D}' which is a normal subgroup of \mathfrak{H} , then the group \mathfrak{G} is not simple.*

Theorem B. [5] *If in the finite group $\mathfrak{G} = \mathfrak{H}\mathfrak{K}$ the orders of the factors \mathfrak{H} and \mathfrak{K} are relatively prime, then every normal subgroup \mathfrak{G} of \mathfrak{G} is of the form $\mathfrak{G} = \mathfrak{H}\mathfrak{K}$ where \mathfrak{H} and \mathfrak{K} are normal subgroups of \mathfrak{H} and \mathfrak{K} respectively.*

After these preliminaries we turn our attention to the proofs of Theorems 1 and 2.

Proof of Theorem 1. If $\mathfrak{D} \neq E$ then the statement is evident.

If $\mathfrak{D} = E$ then each element of \mathfrak{G} may be represented uniquely in the form HK and also in the form $K'H'$ (where $H, H' \in \mathfrak{H}$ and $K, K' \in \mathfrak{K}$). Therefore in $HK = K'H'$ the elements H' and K' are uniquely determined for any given H and K . If K is fixed and H runs over all elements of \mathfrak{H} , then H' also runs over all elements of \mathfrak{H} . The system Σ_K of the corresponding solutions K' of the equation $HK = K'H'$ (each K' taken as many times as it is a solution) is uniquely defined by K .

Let the order of \mathfrak{H} be not less than the order of \mathfrak{K} . Then for every K , the system Σ_K contains K at least twice, in fact, if $K = E$, then the statement is evident. If $K \neq E$, then the system Σ_K does not contain the unit element, thus the statement also holds. The elements H and H' for which the relation $HK = KH'$ holds, form two groups \mathfrak{M} and \mathfrak{M}' respectively. Clearly, $K^{-1}\mathfrak{M}K = \mathfrak{M}$. If all Σ_K contain no elements other than K , then \mathfrak{H} is a normal subgroup of \mathfrak{G} . If $K \neq \bar{K} \in \Sigma_K$ then $\bar{K}^{-1}\mathfrak{M}\bar{K} = \mathfrak{M}$; in fact, if $K^{-1}\mathfrak{M}K = \mathfrak{M}'$ and $\bar{H}K = \bar{K}\bar{H}'$ then $\bar{H}^{-1}\bar{K}^{-1}\bar{H}\mathfrak{M}\bar{H}^{-1}\bar{K}\bar{H}' = \mathfrak{M}$, i. e. $\bar{K}^{-1}\mathfrak{M}\bar{K} = \mathfrak{M}'$ since \mathfrak{H} is an abelian group. For the element $K\bar{K}^{-1} = K^*$ we have $K^{*-1}\mathfrak{M}K^* = \mathfrak{M}$ thus every element of Σ_{K^*} transforms the group \mathfrak{M} into itself.

Therefore if $\{\Sigma_{K^*}\} = \mathfrak{K}$, then \mathfrak{M} is a normal subgroup of \mathfrak{G} .

If $\{\Sigma_{K^*}\} = \mathfrak{K}' \subset \mathfrak{K}$, then the elements $\mathfrak{H}\mathfrak{K}'$ form a proper subgroup of \mathfrak{G} . This will be proved in two steps:

1) $H_i K^* = K_i^* H_i$ implies $\Sigma_{K^*} = \Sigma_{K_i^*}$ ($i = 1, 2, \dots, h$). For, if in $H_i K^* = K_i^* H_i$ ($i = 1, 2, \dots, h$ where h is the order of \mathfrak{H}) we substitute K^* by $K^* = H_k^{-1} K_k^* H_k$ (k is fixed) the statement follows.

2) $HK = K'H'$ and $H'\bar{K} = \bar{K}'H''$ ($K, K', \bar{K}, \bar{K}' \in \Sigma_{K^*}$) imply that with $K\bar{K}$ also $K'\bar{K}'$ belongs to \mathfrak{K}' .

From 1) and 2) it follows that $\mathfrak{H}\mathfrak{K}' = \mathfrak{K}'\mathfrak{H}$, i. e. the product of \mathfrak{H} and \mathfrak{K}' is a subgroup of \mathfrak{G} . Therefore \mathfrak{G} is factorisable by $\mathfrak{H}\mathfrak{K}'$ and \mathfrak{K} thus by Theorem A the group \mathfrak{G} is not simple.

Proof of Theorem 2. (By induction.) The statement is known to hold for groups of order $p \cdot q$ with different primes p and q . Let the order of \mathfrak{G} be n and suppose the theorem true for factorisable groups of order $< n$. Let $\overline{\mathfrak{G}}$ be a proper normal subgroup of \mathfrak{G} ; the existence of such a subgroup follows from Theorem 1. From Theorem B we may conclude that $\overline{\mathfrak{G}} = \overline{\mathfrak{M}}\overline{\mathfrak{N}}$ where $\overline{\mathfrak{M}}$ and $\overline{\mathfrak{N}}$ are normal subgroups of $\overline{\mathfrak{G}}$ and \mathfrak{K} respectively. The groups $\mathfrak{H}\overline{\mathfrak{M}}\mathfrak{N} = \mathfrak{H}\overline{\mathfrak{N}} = \mathfrak{H}$ and $\mathfrak{K}\overline{\mathfrak{M}}\mathfrak{N} = \mathfrak{K}\overline{\mathfrak{N}} = \mathfrak{K}$ are subgroups of \mathfrak{G} and clearly at least one of them is proper. As is easily seen, $\overline{\mathfrak{G}}/\overline{\mathfrak{M}}\mathfrak{N} = \overline{\mathfrak{H}}/\overline{\mathfrak{M}}\mathfrak{N} \cdot \mathfrak{K}/\overline{\mathfrak{M}}\mathfrak{N}$ and $\overline{\mathfrak{H}}/\overline{\mathfrak{M}}\mathfrak{N}$ as well as $\mathfrak{K}/\overline{\mathfrak{M}}\mathfrak{N}$ are abelian groups. The orders of the factor-groups $\overline{\mathfrak{H}}/\overline{\mathfrak{M}}\mathfrak{N}$ and $\mathfrak{K}/\overline{\mathfrak{M}}\mathfrak{N}$ are relatively prime; consequently, by the induction hypothesis, $\overline{\mathfrak{G}}/\overline{\mathfrak{M}}\mathfrak{N}$ has a normal subgroup of prime index p . Therefore the group \mathfrak{G} has a normal subgroup of index p ; this normal subgroup is factorisable by two of its proper abelian subgroups of relatively prime orders (Theorem B), or it is abelian and so the proof is completed.

References.

G. CASADIO

- [1] Costruzione di gruppi come prodotto di sottogruppi permutabili, *Rendiconti di mat. e delle sue applicazioni*, (V) 2 (1941), fasc. III—IV.

L. RÉDEI

- [2] Anwendungen des schiefen Produktes in der Gruppentheorie, *Journal für die reine und angewandte Math.*, 188 (1951), pp. 201—227.
[3] Zur Theorie der faktorisierten Gruppen, *Acta Math. Academiae Scientiarum Hungaricae*. (Under press.)

J. SZÉP

- [4] Über die als Produkt zweier Untergruppen darstellbaren endlichen Gruppen, *Commentarii Math. Helvetici*, 22 (1948), pp. 31—33.
[5] On the structure of groups which can be represented as the product of two subgroups, *these Acta*, 12 A (1950), pp. 57—61.
[6] On simple groups, *Publicationes Math., Debrecen*, 1 (1949), p. 98.
[7] On factorisable simple groups. (To be published in *these Acta*.)

J. SZÉP—L. RÉDEI

- [8] On factorisable groups, *these Acta*, 13 (1950), pp. 235—238.

G. ZAPPA

- [9] Costruzione dei gruppi prodotto di due dati sottogruppi permutabili tra loro, *Atti Secondo Congresso Unione Mat. Italiana Bologna 1940*, pp. 115—125.

(Received April 1, 1950.)